



中山大學
SUN YAT-SEN UNIVERSITY

Bose-Hubbard Model

學大山中立國

中山大学物理学院



报告人：路尚润

2024.4.7

目录

CONTENTS

01

Cold Bosonic Atoms in Optical Lattices

02

Ultracold atoms in optical lattices
generated by quantized light fields

03

Quantum phases in an optical lattice



中山大學
SUN YAT-SEN UNIVERSITY

Part.01

Cold Bosonic Atoms in Optical Lattices

• Hamiltonian

Hamilton operator for bosonic atoms

$$H = \int d^3x \psi^\dagger(\mathbf{x}) \left(-\frac{\hbar^2}{2m} \nabla^2 + V_0(\mathbf{x}) + V_T(\mathbf{x}) \right) \psi(\mathbf{x}) + \frac{1}{2} \frac{4\pi a_s \hbar^2}{m} \int d^3x \psi^\dagger(\mathbf{x}) \psi^\dagger(\mathbf{x}) \psi(\mathbf{x}) \psi(\mathbf{x}),$$

$\psi(\mathbf{x})$ is a boson field operator for atoms in a given internal atomic state

$V_0(x)$ is the optical lattice potential

$V_T(x)$ describes an additional (slowly varying) external trapping potential

$$V_0(\mathbf{x}) = \sum_{j=1}^3 V_{j0} \sin^2(kx_j)$$

$$k = 2\pi/\lambda \quad a = \lambda/2$$

Expanding the field operators in the Wannier basis and keeping only the lowest vibrational states

$$\psi(\mathbf{x}) = \sum_i b_i w(\mathbf{x} - \mathbf{x}_i)$$

$$\hat{\Psi}(\mathbf{r}) = \sum_j \hat{b}_j \phi_j(\mathbf{r})$$

$$H = -J \sum_{\langle i,j \rangle} b_i^\dagger b_j + \sum_i \epsilon_i \hat{n}_i + \frac{1}{2} U \sum_i \hat{n}_i (\hat{n}_i - 1),$$

$$\Psi_\ell(r_\ell) = (k_0^2 \tilde{\omega}_\ell / \pi)^{1/4} e^{-\tilde{\omega}_\ell k_0^2 r_\ell^2 / 2}$$

近邻交叠积分最大

• Tight-Binding Model

$$\psi(\mathbf{x}) = \sum_i b_i w(\mathbf{x} - \mathbf{x}_i)$$

$$H = -J \sum_{\langle i,j \rangle} b_i^\dagger b_j + \sum_i \epsilon_i \hat{n}_i + \frac{1}{2} U \sum_i \hat{n}_i (\hat{n}_i - 1)$$


where the operators $\hat{n}_i = b_i^\dagger b_i$ count the number of bosonic atoms at lattice site i ;

the annihilation and creation operators b_i and b_i^\dagger obey the canonical commutation relations $[b_i, b_j^\dagger] = \delta_{ij}$.

The parameters $U = 4\pi a_s \hbar^2 \int d^3x |w(\mathbf{x})|^4 / m$ correspond to the strength of the on site repulsion of two atoms on the lattice site i .

$J = - \int d^3x w^*(\mathbf{x} - \mathbf{x}_i) \left[-\frac{\hbar^2}{2m} \nabla^2 + V_0(\mathbf{x}) \right] w(\mathbf{x} - \mathbf{x}_j)$ is the hopping matrix element between adjacent sites i, j .

$\epsilon_i = \int d^3x V_T(\mathbf{x}) |w(\mathbf{x} - \mathbf{x}_i)|^2 \approx V_T(\mathbf{x}_i)$ describes an energy offset of each lattice site.





中山大學
SUN YAT-SEN UNIVERSITY

Part.02

Ultracold atoms in optical lattices generated by quantized light fields

• Single Atom Hamiltonian

N个质量为m，跃迁频率为 ω_{eg} 的原子，与频率 ω_c 的驻波单模腔场相互作用，还有一个 ω_p 频率、与原子最大耦合强度为 h_0 的振幅为 η 的相干驱动光，弱耦合采用RWA，得到JC Model。

$$H^{(1)} = H_A^{(1)} + H_R^{(1)} + H_{Int}^{(1)}.$$

$$H_A^{(1)} = \frac{\hat{\mathbf{p}}^2}{2m} + V_e(\mathbf{x})\sigma^+\sigma^- + V_g(\mathbf{x})\sigma^-\sigma^+ + \hbar\omega_{eg}\sigma^+\sigma^- - i\hbar h(\mathbf{x}) (\sigma^+ e^{-i\omega_p t} - \sigma^- e^{i\omega_p t})$$

$$H_R^{(1)} = \hbar\omega_c a^\dagger a - i\hbar\eta (a e^{i\omega_p t} - a^\dagger e^{-i\omega_p t}),$$

$$H_{Int}^{(1)} = -i\hbar g(\mathbf{x}) (\sigma^+ a - \sigma^- a^\dagger).$$




$$U(t) = \exp[i\omega_p t (\sigma^+\sigma^- + a^\dagger a)]$$

$$H_A^{(1)} = \frac{\hat{\mathbf{p}}^2}{2m} + V_e(\mathbf{x})\sigma^+\sigma^- + V_g(\mathbf{x})\sigma^-\sigma^+ - \hbar\Delta_a\sigma^+\sigma^- - i\hbar h(\mathbf{x}) (\sigma^+ - \sigma^-)$$

$$H_R^{(1)} = -\hbar\Delta_c a^\dagger a - i\hbar\eta (a - a^\dagger),$$

$$H_{Int}^{(1)} = -i\hbar g(\mathbf{x}) (\sigma^+ a - \sigma^- a^\dagger),$$



detuning

$$\Delta_c = \omega_p - \omega_c, \Delta_a = \omega_p - \omega_{eg}$$

• Second Quantization

$$H_A^{(1)} = \frac{\hat{\mathbf{p}}^2}{2m} + V_e(\mathbf{x})\sigma^+\sigma^- + V_g(\mathbf{x})\sigma^-\sigma^+ - \hbar\Delta_a\sigma^+\sigma^- - i\hbar h(\mathbf{x})(\sigma^+ - \sigma^-)$$

$$H_R^{(1)} = -\hbar\Delta_c a^\dagger a - i\hbar\eta(a - a^\dagger),$$

$$H_{Int}^{(1)} = -i\hbar g(\mathbf{x})(\sigma^+ a - \sigma^- a^\dagger),$$




$$H = H_A + H_R + H_{A-R} + H_{A-P} + H_{A-A}.$$

$$H_A = \int d^3\mathbf{x} \left[\Psi_g^\dagger(\mathbf{x}) \left(-\frac{\hbar^2}{2m} \nabla^2 + V_g(\mathbf{x}) \right) \Psi_g(\mathbf{x}) + \Psi_e^\dagger(\mathbf{x}) \left(-\frac{\hbar^2}{2m} \nabla^2 - \hbar\Delta_a + V_e(\mathbf{x}) \right) \Psi_e(\mathbf{x}) \right]$$

$$H_{A-A} = \frac{U}{2} \int d^3\mathbf{x} \Psi_g^\dagger(\mathbf{x}) \Psi_g^\dagger(\mathbf{x}) \Psi_g(\mathbf{x}) \Psi_g(\mathbf{x}), \quad U = 4\pi a_s \hbar^2 / m.$$

$$H_{A-R} = -i\hbar \int d^3\mathbf{x} \Psi_g^\dagger(\mathbf{x}) g(\mathbf{x}) a^\dagger \Psi_e(\mathbf{x}) + \text{h.c.},$$

$$H_{A-P} = -i\hbar \int d^3\mathbf{x} \Psi_g^\dagger(\mathbf{x}) h(\mathbf{x}) \Psi_e(\mathbf{x}) + \text{h.c..}$$



$$\hat{\Psi}_e^\dagger(\mathbf{r}) \hat{\Psi}_g(\mathbf{r}) = \hat{\sigma}_+(\mathbf{r})$$

$$\hat{\Psi}_g^\dagger(\mathbf{r}) \hat{\Psi}_e(\mathbf{r}) = \hat{\sigma}_-(\mathbf{r})$$

$$[\Psi_f(\mathbf{x}), \Psi_{f'}^\dagger(\mathbf{x}')] = \delta^3(\mathbf{x} - \mathbf{x}') \delta_{f,f'}$$

$$[\Psi_f(\mathbf{x}), \Psi_{f'}(\mathbf{x}')] = [\Psi_f^\dagger(\mathbf{x}), \Psi_{f'}^\dagger(\mathbf{x}')]=0,$$

• Heisenberg Equations

$$\frac{\partial \Psi_e(\mathbf{x})}{\partial t} = i \left[\frac{\hbar}{2m} \nabla^2 - \frac{V_e(\mathbf{x})}{\hbar} + \Delta_a \right] \Psi_e(\mathbf{x}) \quad \text{free evolution}$$

+ $[g(\mathbf{x})a + h(\mathbf{x})] \Psi_g(\mathbf{x})$. 光子吸收及基态湮灭

$$\frac{\partial \Psi_g(\mathbf{x})}{\partial t} = i \left[\frac{\hbar}{2m} \nabla^2 - \frac{V_g(\mathbf{x})}{\hbar} - \frac{U}{\hbar} \Psi_g^\dagger(\mathbf{x}) \Psi_g(\mathbf{x}) \right] \Psi_g(\mathbf{x})$$

± $[g(\mathbf{x})a^\dagger + h(\mathbf{x})] \Psi_e(\mathbf{x})$. 光子发射及激发态产生

$$\frac{\partial a}{\partial t} = i \Delta_c a + \eta \pm \int d^3 \mathbf{x} g(\mathbf{x}) \Psi_g^\dagger(\mathbf{x}) \Psi_e(\mathbf{x}).$$

T=0, 弱激发态



large atom-pump detuning Δ_a

$$\Psi_e(\mathbf{x}, t) = \pm \frac{i}{\Delta_a} [h(\mathbf{x}) + g(\mathbf{x})a(t)] \Psi_g(\mathbf{x}, t). \quad \frac{\partial a}{\partial t} = i \left[\Delta_c - \frac{1}{\Delta_a} \int d^3 \mathbf{x} g^2(\mathbf{x}) \Psi_g^\dagger(\mathbf{x}) \Psi_g(\mathbf{x}) \right] a - \frac{i}{\Delta_a} \int d^3 \mathbf{x} h(\mathbf{x}) \Psi_g^\dagger(\mathbf{x}) \Psi_g(\mathbf{x}) + \eta.$$

$$\frac{\partial \Psi_g(\mathbf{x})}{\partial t} = i \left[\frac{\hbar}{2m} \nabla^2 - \frac{V_g(\mathbf{x})}{\hbar} - \frac{h^2(\mathbf{x})}{\Delta_a} - \frac{g^2(\mathbf{x})}{\Delta_a} a^\dagger a - \frac{h(\mathbf{x})g(\mathbf{x})}{\Delta_a} (a + a^\dagger) - \frac{U}{\hbar} \Psi_g^\dagger(\mathbf{x}) \Psi_g(\mathbf{x}) \right] \Psi_g(\mathbf{x})$$

$$\hat{\Psi}_e^\dagger(\mathbf{r}) \hat{\Psi}_g(\mathbf{r}) = \hat{\sigma}_+(\mathbf{r})$$

$$\hat{\Psi}_g^\dagger(\mathbf{r}) \hat{\Psi}_e(\mathbf{r}) = \hat{\sigma}_-(\mathbf{r})$$

$$[\Psi_f(\mathbf{x}), \Psi_{f'}^\dagger(\mathbf{x}')] = \delta^3(\mathbf{x} - \mathbf{x}') \delta_{f,f'}$$

$$[\Psi_f(\mathbf{x}), \Psi_{f'}(\mathbf{x}')] = [\Psi_f^\dagger(\mathbf{x}), \Psi_{f'}^\dagger(\mathbf{x}')] = 0,$$

$g(\mathbf{x})$

• Effective Hamiltonian

$$\frac{\partial a}{\partial t} = i \left[\Delta_c - \frac{1}{\Delta_a} \int d^3x g^2(\mathbf{x}) \Psi_g^\dagger(\mathbf{x}) \Psi_g(\mathbf{x}) \right] a - \frac{i}{\Delta_a} \int d^3x h(\mathbf{x}) \Psi_g^\dagger(\mathbf{x}) \Psi_g(\mathbf{x}) + \eta.$$

$$\frac{\partial \Psi_g(\mathbf{x})}{\partial t} = i \left[\frac{\hbar}{2m} \nabla^2 - \frac{V_g(\mathbf{x})}{\hbar} - \frac{h^2(\mathbf{x})}{\Delta_a} - \frac{g^2(\mathbf{x})}{\Delta_a} a^\dagger a - \frac{h(\mathbf{x})g(\mathbf{x})}{\Delta_a} (a + a^\dagger) - \frac{U}{\hbar} \Psi_g^\dagger(\mathbf{x}) \Psi_g(\mathbf{x}) \right] \Psi_g(\mathbf{x})$$

$g(\mathbf{x})$



$$H_{\text{eff}} = \int d^3x \Psi_g^\dagger(\mathbf{x}) \left\{ -\frac{\hbar^2}{2m} \nabla^2 + V_g(\mathbf{x}) + \frac{\hbar}{\Delta_a} [h^2(\mathbf{x}) + g^2(\mathbf{x}) a^\dagger a + h(\mathbf{x})g(\mathbf{x}) (a + a^\dagger)] \right\} \Psi_g(\mathbf{x}) \\ + \frac{U}{2} \int d^3x \Psi_g^\dagger(\mathbf{x}) \Psi_g^\dagger(\mathbf{x}) \Psi_g(\mathbf{x}) \Psi_g(\mathbf{x}) - i\hbar\eta (a - a^\dagger) - \hbar\Delta_c a^\dagger a.$$



Single particle Hamiltonian

$$H_{\text{eff}}^{(1)} = \frac{\mathbf{p}^2}{2m} + V_g(\mathbf{x}) + \frac{\hbar}{\Delta_a} [h^2(\mathbf{x}) + g^2(\mathbf{x}) a^\dagger a + h(\mathbf{x})g(\mathbf{x}) (a + a^\dagger)] - i\hbar\eta (a - a^\dagger) - \hbar\Delta_c a^\dagger a.$$



• 1D Open System

$$\begin{aligned} h(\mathbf{x}) &= h_0 \cos(k_p y) & \dot{\varrho} &= \frac{1}{i\hbar} [H_{\text{eff}}, \varrho] + \mathcal{L}\varrho. \\ V_g(\mathbf{x}) &= V_g(x) & & \text{Cavity loss } \kappa \\ g(x) &= g_0 \cos(kx) & \mathcal{L}\varrho &= \kappa (2a\varrho a^\dagger - a^\dagger a\varrho - \varrho a^\dagger a) \end{aligned}$$

$$\frac{\partial a}{\partial t} = i \left[\Delta_c - \frac{1}{\Delta_a} \int d^3 \mathbf{x} g^2(\mathbf{x}) \Psi_g^\dagger(\mathbf{x}) \Psi_g(\mathbf{x}) \right] a - \frac{i}{\Delta_a} \int d^3 \mathbf{x} h(\mathbf{x}) \Psi_g^\dagger(\mathbf{x}) \Psi_g(\mathbf{x}) + \eta.$$



Langevin equation

$$\dot{a} = \left\{ i \left[\Delta_c - \frac{g_0^2}{\Delta_a} \int dx \Psi_g^\dagger(x) \cos^2(kx) \Psi_g(x) \right] - \kappa \right\} a - i \frac{g_0 h_0}{\Delta_a} \int dx \Psi_g^\dagger(x) \cos(kx) \Psi_g(x) + \eta + \Gamma_{in}$$

Cavity loss κ

Noise operator

外部真空 $T=0$, 噪声算符的
平均值为0, 不进入动力学

•Bose-Hubbard Hamiltonian

$$\Psi_g(x) = \sum_k b_k w(x - x_k) \quad w(x) = w_0(x)$$

$$H_{\text{eff}} = \int d^3x \Psi_g^\dagger(\mathbf{x}) \left\{ -\frac{\hbar^2}{2m} \nabla^2 + V_g(\mathbf{x}) + \frac{\hbar}{\Delta_a} [h^2(\mathbf{x}) + g^2(\mathbf{x}) a^\dagger a + h(\mathbf{x}) g(\mathbf{x}) (a + a^\dagger)] \right\} \Psi_g(\mathbf{x}) \\ + \frac{U}{2} \int d^3x \Psi_g^\dagger(\mathbf{x}) \Psi_g^\dagger(\mathbf{x}) \Psi_g(\mathbf{x}) \Psi_g(\mathbf{x}) - i\hbar\eta (a - a^\dagger) - \hbar\Delta_c a^\dagger a.$$

$$V_g(x) = V_{\text{cl}} \cos^2(k_F x)$$



低激发腔场势阱太小，故增加外部捕获势

$$H = \sum_{k,l} E_{kl} b_k^\dagger b_l + (\hbar U_0 a^\dagger a + V_{\text{cl}}) \sum_{k,l} J_{kl} b_k^\dagger b_l \\ + \hbar\eta_{\text{eff}} (a + a^\dagger) \sum_{k,l} \tilde{J}_{kl} b_k^\dagger b_l - i\hbar\eta (a - a^\dagger) \\ + \frac{1}{2} \sum_{i,j,k,l} U_{ijkl} b_i^\dagger b_j^\dagger b_k b_l - \hbar\Delta_c a^\dagger a,$$

$$E_{kl} = \int dx w(x - x_k) \left(-\frac{\hbar^2}{2m} \nabla^2 \right) w(x - x_l), \quad \text{cos}(k) \text{在两个井之间一正一负} \\ J_{kl} = \int dx w(x - x_k) \cos^2(kx) w(x - x_l), \\ \tilde{J}_{kl} = \int dx w(x - x_k) \cos(kx) w(x - x_l). \quad \tilde{J}_{k,k+1} = 0 \\ U_{ijkl} = g_{1D} \int dx w(x - x_i) w(x - x_j) w(x - x_k) w(x - x_l)$$



中山大學
SUN YAT-SEN UNIVERSITY

Part.03


Quantum phases in an optical lattice



•Bogoliubov Approximation

$$H = -t \sum_{\langle i,j \rangle} c_i^\dagger c_j + \frac{1}{2} U \sum_i c_i^\dagger c_i^\dagger c_i c_i - \mu \sum_i c_i^\dagger c_i,$$

momentum space



$$c_i = \frac{1}{\sqrt{N_s}} \sum_{\mathbf{k}} a_{\mathbf{k}} e^{-i\mathbf{k} \cdot \mathbf{r}_i}, \quad c_i^\dagger = \frac{1}{\sqrt{N_s}} \sum_{\mathbf{k}} a_{\mathbf{k}}^\dagger e^{i\mathbf{k} \cdot \mathbf{r}_i},$$

$$\sum_i e^{-i(\mathbf{k}-\mathbf{k}') \cdot \mathbf{r}_i} = N_s \delta_{\mathbf{k}, \mathbf{k}'}$$

$$H = \sum_{\mathbf{k}} (-\bar{\epsilon}_{\mathbf{k}} - \mu) a_{\mathbf{k}}^\dagger a_{\mathbf{k}} + \frac{1}{2} \frac{U}{N_s} \sum_{\mathbf{k}} \sum_{\mathbf{k}'} \sum_{\mathbf{k}''} \sum_{\mathbf{k}'''} a_{\mathbf{k}}^\dagger a_{\mathbf{k}'}^\dagger a_{\mathbf{k}''} a_{\mathbf{k}'''} \delta_{\mathbf{k}+\mathbf{k}', \mathbf{k}''+\mathbf{k}'''},$$


$$\bar{\epsilon}_{\mathbf{k}} = 2t \sum_{j=1}^d \cos(k_j a)$$

For a Bose condensed gas $N_0 \gg 1$: $N_0 = \langle a_0^\dagger a_0 \rangle \approx \langle a_0 a_0^\dagger \rangle$ $N_0 = \langle a_0^\dagger \rangle \langle a_0 \rangle$ $\langle a_0^\dagger \rangle = \langle a_0 \rangle = \sqrt{N_0}$

Bogoliubov approach:

$$a_0^\dagger \rightarrow \sqrt{N_0} + a_0^\dagger, \quad a_0 \rightarrow \sqrt{N_0} + a_0,$$

minimizing the energy of the gas with respect to the number of condensate atoms N_0



•Bogoliubov Approximation

波动线性项=0 $H^{(1)} = \left(-\bar{\epsilon}_0 - \mu + \frac{U}{N_s} N_0 \right) \sqrt{N_0} (a_0^\dagger + a_0)$,



$\mu = Un_0 - zt$ condensate density $n_0 = N_0 / N_s$ the number of nearest neighbors $z = 2d$

化学势能是增加一个粒子所需的能量：包含在自身格点的相互作用能增加和可能hopping的能量减小

Effective Hamiltonian

$$H^{\text{eff}} = \left(-zt - \mu + \frac{1}{2} Un_0 \right) N_0 + \sum_{\mathbf{k}} (-\bar{\epsilon}_{\mathbf{k}} - \mu) a_{\mathbf{k}}^\dagger a_{\mathbf{k}} + \frac{1}{2} Un_0 \sum_{\mathbf{k}} (a_{\mathbf{k}} a_{-\mathbf{k}} + 4a_{\mathbf{k}}^\dagger a_{\mathbf{k}} + a_{-\mathbf{k}}^\dagger a_{-\mathbf{k}}).$$

Matrix form 令 $\epsilon_{\mathbf{k}} = zt - \bar{\epsilon}_{\mathbf{k}}$

$$H^{\text{eff}} = -\frac{1}{2} Un_0 N_0 - \frac{1}{2} \sum_{\mathbf{k}} (\epsilon_{\mathbf{k}} + Un_0) + \frac{1}{2} \sum_{\mathbf{k}} (a_{\mathbf{k}}^\dagger, a_{-\mathbf{k}}) \times \begin{bmatrix} \epsilon_{\mathbf{k}} + Un_0 & Un_0 \\ Un_0 & \epsilon_{\mathbf{k}} + Un_0 \end{bmatrix} \begin{pmatrix} a_{\mathbf{k}} \\ a_{-\mathbf{k}}^\dagger \end{pmatrix}$$

$$H = \sum_{\mathbf{k}} (-\bar{\epsilon}_{\mathbf{k}} - \mu) a_{\mathbf{k}}^\dagger a_{\mathbf{k}} + \frac{1}{2} \frac{U}{N_s} \sum_{\mathbf{k}} \sum_{\mathbf{k}'} \sum_{\mathbf{k}''} \sum_{\mathbf{k}'''} a_{\mathbf{k}}^\dagger a_{\mathbf{k}'}^\dagger a_{\mathbf{k}''} a_{\mathbf{k}'''} \delta_{\mathbf{k} + \mathbf{k}', \mathbf{k}'' + \mathbf{k}'''},$$

$a_0^\dagger \rightarrow \sqrt{N_0} + a_0^\dagger, \quad a_0 \rightarrow \sqrt{N_0} + a_0,$ minimizing the energy



•Bogoliubov Approximation

$$H^{\text{eff}} = -\frac{1}{2} Un_0 N_0 - \frac{1}{2} \sum_{\mathbf{k}} (\epsilon_{\mathbf{k}} + Un_0) + \frac{1}{2} \sum_{\mathbf{k}} (a_{\mathbf{k}}^\dagger, a_{-\mathbf{k}}) \times \begin{bmatrix} \epsilon_{\mathbf{k}} + Un_0 & Un_0 \\ Un_0 & \epsilon_{\mathbf{k}} + Un_0 \end{bmatrix} \begin{pmatrix} a_{\mathbf{k}} \\ a_{-\mathbf{k}}^\dagger \end{pmatrix}$$

diagonalized

$$\begin{pmatrix} b_{\mathbf{k}} \\ b_{-\mathbf{k}}^\dagger \end{pmatrix} = \begin{bmatrix} u_{\mathbf{k}} & v_{\mathbf{k}} \\ v_{\mathbf{k}}^* & u_{\mathbf{k}}^* \end{bmatrix} \begin{pmatrix} a_{\mathbf{k}} \\ a_{-\mathbf{k}}^\dagger \end{pmatrix} \equiv \mathbf{B} \begin{pmatrix} a_{\mathbf{k}} \\ a_{-\mathbf{k}}^\dagger \end{pmatrix}$$

$$|u_{\mathbf{k}}|^2 - |v_{\mathbf{k}}|^2 = 1$$

$$n = \frac{1}{N_s} \sum_{\mathbf{k}} \langle a_{\mathbf{k}}^\dagger a_{\mathbf{k}} \rangle_{H^{\text{eff}}},$$

$$\hbar \omega_{\mathbf{k}} = \sqrt{\epsilon_{\mathbf{k}}^2 + 2 Un_0 \epsilon_{\mathbf{k}}} \quad \rightarrow \quad |v_{\mathbf{k}}|^2 = |u_{\mathbf{k}}|^2 - 1 = \frac{1}{2} \left(\frac{\epsilon_{\mathbf{k}} + Un_0}{\hbar \omega_{\mathbf{k}}} - 1 \right).$$

$$H^{\text{eff}} = -\frac{1}{2} Un_0 N_0 + \frac{1}{2} \sum_{\mathbf{k}} [\hbar \omega_{\mathbf{k}} - (\epsilon_{\mathbf{k}} + Un_0)] + \sum_{\mathbf{k}} \hbar \omega_{\mathbf{k}} b_{\mathbf{k}}^\dagger b_{\mathbf{k}}$$

$$n = n_0 + \frac{1}{N_s} \sum_{\mathbf{k} \neq 0} [(|u_{\mathbf{k}}|^2 + |v_{\mathbf{k}}|^2) \langle b_{\mathbf{k}}^\dagger b_{\mathbf{k}} \rangle_{H^{\text{eff}}} + |v_{\mathbf{k}}|^2].$$


Bose distribution

$$n = n_0 + \frac{1}{N_s} \sum_{\mathbf{k} \neq 0} \left(\frac{\epsilon_{\mathbf{k}} + Un_0}{\hbar \omega_{\mathbf{k}}} \frac{1}{e^{\beta \hbar \omega_{\mathbf{k}} - 1}} + \frac{\epsilon_{\mathbf{k}} + Un_0 - \hbar \omega_{\mathbf{k}}}{2 \hbar \omega_{\mathbf{k}}} \right) \quad \begin{array}{c} T=0, \beta \rightarrow \infty \\ k = 2\pi q/a \end{array} \rightarrow$$

$$n = n_0 + \frac{1}{2} \int_{-1/2}^{1/2} d\mathbf{q} \left(\frac{\epsilon_{\mathbf{q}} + Un_0}{\hbar \omega_{\mathbf{q}}} - 1 \right)$$

$$\epsilon_{\mathbf{q}} = 2 t \sum_{j=1}^d [1 - \cos(2\pi q_j)]$$


$$\hbar \omega_{\mathbf{q}} = (\epsilon_{\mathbf{q}}^2 + 2 Un_0 \epsilon_{\mathbf{q}})^{1/2}$$



$$|\Psi\rangle_{J/U=0} \propto \prod_i |n_i\rangle$$



$$|\Psi\rangle_{U/J=0} \propto \left(\sum_i \hat{b}_i^\dagger \right)^N |0\rangle$$



Probing correlated quantum many-body systems at the single-particle level
-Manuel Endres



•Bogoliubov Approximation

$$n = n_0 + \frac{1}{2} \int_{-1/2}^{1/2} d\mathbf{q} \left(\frac{\epsilon_{\mathbf{q}} + Un_0}{\hbar \omega_{\mathbf{q}}} - 1 \right)$$

对于 $n=Z$, 应当没有超流解

Asymptotic behavior $U/t \rightarrow \infty$ $\epsilon_{\mathbf{q}} \leq 4\pi^2 |\mathbf{q}|^2 t$.


$$\int_{-1/2}^{1/2} d\mathbf{q} \frac{\epsilon_{\mathbf{q}} + Un_0}{\sqrt{\epsilon_{\mathbf{q}}^2 + 2U\epsilon_{\mathbf{q}}n_0}} \geq \frac{1}{2\pi} \sqrt{\frac{Un_0}{2t}} \int_{-1/2}^{1/2} \frac{d\mathbf{q}}{|\mathbf{q}|} = I_d$$

$$n \approx n_0 + \frac{1}{4\pi} \sqrt{\frac{Un_0}{2t}} I_d - \frac{1}{2}$$

\rightarrow

$$n_0 = \left(\frac{1}{2} \sqrt{\frac{I_d^2}{16\pi^2} \frac{U}{2t} + 4n + 2} - \frac{I_d}{8\pi} \sqrt{\frac{U}{2t}} \right)^2$$

一直存在正的解，没有超流-绝缘体转变，只有极限才有 $n_0=0$



- Mean-field Approach

The superfluid order parameter $\psi = \sqrt{n_i} = \langle c_i^\dagger \rangle = \langle c_i \rangle$

$$H = -t \sum_{\langle i,j \rangle} c_i^\dagger c_j + \frac{1}{2} U \sum_i c_i^\dagger c_i^\dagger c_i c_i - \mu \sum_i c_i^\dagger c_i,$$

做代换

$$c_i^\dagger c_j = \langle c_i^\dagger \rangle c_j + c_i^\dagger \langle c_j \rangle - \langle c_i^\dagger \rangle \langle c_j \rangle = \psi(c_i^\dagger + c_j) - \psi^2$$



$$H^{\text{eff}} = -zt\psi \sum_i (c_i^\dagger + c_i) + zt\psi^2 N_s + \frac{1}{2} U \sum_i c_i^\dagger c_i^\dagger c_i c_i - \mu \sum_i c_i^\dagger c_i,$$

$$\bar{U} = U/zt, \bar{\mu} = \mu/zt$$



消除维度依赖

the number of nearest neighbors $z=2d$

$$H_i^{\text{eff}} = \frac{1}{2} \bar{U} \hat{n}_i (\hat{n}_i - 1) - \bar{\mu} \hat{n}_i - \psi(c_i^\dagger + c_i) + \psi^2$$

转化为最小化单粒子基态能量

- Mean-field Approach

Second-order perturbation theory $H^{\text{eff}} = H^{(0)} + \psi V$

$$H^{(0)} = \frac{1}{2} \bar{U} \hat{n}(\hat{n}-1) - \bar{\mu} \hat{n} + \psi^2, \quad V = -(c^\dagger + c), \quad E_g^{(0)} = \{E_n^{(0)} | n=0,1,2,\dots\}_{\min}.$$

$$E_g^{(0)} = \begin{cases} 0 & \text{if } \bar{\mu} < 0, \\ \frac{1}{2} \bar{U} g(g-1) - \bar{\mu} g & \text{if } \bar{U}(g-1) < \bar{\mu} < \bar{U}g. \end{cases}$$

比较 g 粒子与 $g \pm 1$ 粒子能量

$$E_g^{(2)} = \psi^2 \sum_{n \neq g} \frac{|\langle g | V | n \rangle|^2}{E_g^{(0)} - E_n^{(0)}}, \quad \xrightarrow{\text{只有两个不为0}} \quad E_g^{(2)} = \frac{g}{\bar{U}(g-1) - \bar{\mu}} + \frac{g+1}{\bar{\mu} - \bar{U}g}.$$

$$E_g(\psi) = a_0(g, \bar{U}, \bar{\mu}) + a_2(g, \bar{U}, \bar{\mu}) \psi^2 + \mathcal{O}(\psi^4), \quad a_2(g, \bar{U}, \bar{\mu}) = \frac{g}{\bar{U}(g-1) - \bar{\mu}} + \frac{g+1}{\bar{\mu} - \bar{U}g} + 1 = 0,$$



- Mean-field Approach

$$E_g(\psi) = a_0(g, \bar{U}, \bar{\mu}) + a_2(g, \bar{U}, \bar{\mu})\psi^2 + \mathcal{O}(\psi^4), \quad a_2(g, \bar{U}, \bar{\mu}) = \frac{g}{U(g-1)-\bar{\mu}} + \frac{g+1}{\bar{\mu}-Ug} + 1 = 0,$$




$a_2 > 0$ 则 $\psi = 0$, $a_2 < 0$ 则 $\psi > 0$, 则 $a_2 = 0$ 为相变点

$$\bar{\mu}_{\pm} = \frac{1}{2} [\bar{U}(2g-1)-1] \pm \frac{1}{2} \sqrt{\bar{U}^2 - 2\bar{U}(2g+1)+1},$$



令两者相等可以得到临界 \bar{U}_c

$$\bar{U}_c = 2g+1 + \sqrt{(2g+1)^2 - 1},$$






- Mean-field Approach


Fourth-order perturbation theory

$$E_g(\psi) = a_0(g, \bar{U}, \bar{\mu}) + a_2(g, \bar{U}, \bar{\mu})\psi^2 + a_4(g, \bar{U}, \bar{\mu})\psi^4,$$

$$a_4(g, \bar{U}, \bar{\mu}) = \frac{g(g-1)}{[\bar{U}(g-1)-\bar{\mu}]^2[\bar{U}(2g-3)-2\bar{\mu}]} + \frac{(g+1)(g+2)}{(\bar{\mu}-\bar{U}g)^2[2\bar{\mu}-\bar{U}(2g+1)]} - \left(\frac{g}{\bar{U}(g-1)-\bar{\mu}} + \frac{g+1}{\bar{\mu}-\bar{U}g} \right) \times \left(\frac{g}{[\bar{U}(g-1)-\bar{\mu}]^2} + \frac{g+1}{(\bar{\mu}-\bar{U}g)^2} \right)$$



$$\bar{U} = 11 \text{ and } \bar{\mu} = 8.9$$



$$\bar{U} = 11 \text{ and } \bar{\mu} = 7.8$$



- Mean-field Approach


$\bar{\mu} = n \bar{U}$ 此时 $n+1$ 和 n 粒子能量相同，利用二重简并微扰(文章写的 n 与 $n-1$)

$$E_g(\psi)|_{\mu=n\bar{U}}=-\frac{1}{2}\bar{U}n(n+1)+\psi^2-|\psi|\sqrt{n+1},$$


grand-canonical ensemble $\psi_{\min}=[-a_2(g,\bar{U},\bar{\mu})/2a_4(g,\bar{U},\bar{\mu})]^{1/2}$

$$n = -\frac{\partial \langle H^{\text{eff}} \rangle}{\partial \mu} = -\frac{\partial E_g(\psi=\psi_{\min})}{\partial \bar{\mu}} = g - \frac{\partial}{\partial \bar{\mu}} \left(\frac{a_2(g,\bar{U},\bar{\mu})^2}{4a_4(g,\bar{U},\bar{\mu})} \right)$$

在 μ_{\pm} 之间， n 为常数！
对应 Mott 绝缘体



$$\bar{U} = U/zt = 11$$



$$\bar{U} = \bar{\mu} = 11$$

• Dispersion Relations

将涨落视为准粒子和准空穴激发，计算二者色散关系

定义复变函数 $a_i^*(\tau)$ and $a_i(\tau)$ $Z = \text{Tr } e^{-\beta \hat{H}} = \int \mathcal{D}a^* \mathcal{D}a \exp\{-S[a^*, a]/\hbar\},$

作用量

$$S[a^*, a] = \int_0^{\hbar\beta} d\tau \left[\sum_i a_i^* \left(\hbar \frac{\partial}{\partial \tau} - \mu \right) a_i - \sum_{ij} t_{ij} a_i^* a_j + \frac{1}{2} U \sum_i a_i^* a_i^* a_i a_i \right],$$

Hubbard-Stratonovich transformation

$$S[a^*, a, \psi^*, \psi] = S[a^*, a] + \int_0^{\hbar\beta} d\tau \sum_{ij} (\psi_i^* - a_i^*) t_{ij} \times (\psi_j - a_j)$$

ψ^* and ψ 是序参量场

$$S[a^*, a, \psi^*, \psi] = \int_0^{\hbar\beta} d\tau \left[\sum_i a_i^* \left(\hbar \frac{\partial}{\partial \tau} - \mu \right) a_i + \frac{1}{2} U \sum_i a_i^* a_i^* a_i a_i - \sum_{ij} t_{ij} (a_i^* \psi_j + \psi_i^* a_j) + \sum_{ij} t_{ij} \psi_i^* \psi_j \right]$$

$S^{(0)}[a^*, a]$ 为 $t_{ij} = 0$ 的项

$$\exp(-S^{\text{eff}}[\psi^*, \psi]/\hbar) \equiv \exp\left(-\frac{1}{\hbar} \int_0^{\hbar\beta} d\tau \sum_{ij} t_{ij} \psi_i^* \psi_j\right) \int \mathcal{D}a^* \mathcal{D}a \times \exp\{-S^{(0)}[a^*, a]/\hbar\} \times \exp\left[-\frac{1}{\hbar} \int_0^{\hbar\beta} d\tau \left(-\sum_{ij} t_{ij} (a_i^* \psi_j + \psi_i^* a_j)\right)\right]$$



• Dispersion Relations

利用Taylor展开到二阶(这里很迷)

$$\begin{aligned} S^{(2)}[\psi^*, \psi] &= -\frac{1}{2\hbar} \left\langle \left(\int_0^{\hbar\beta} d\tau \sum_{ij} t_{ij} (\alpha_i^* \psi_j + \psi_i^* \alpha_j) \right)^2 \right\rangle_{S^{(0)}} + \int_0^{\hbar\beta} d\tau \sum_{ij} t_{ij} \psi_i^* \psi_j \\ &= -\frac{1}{2\hbar} \left\langle \int_0^{\hbar\beta} d\tau \int_0^{\hbar\beta} d\tau' \sum_{iji'j'} t_{ij} t_{i'j'} (\alpha_i^* \psi_j + \psi_i^* \alpha_j) (\alpha_{i'}^* \psi_{j'} + \psi_{i'}^* \alpha_{j'}) \right\rangle_{S^{(0)}} + \int_0^{\hbar\beta} d\tau \sum_{ij} t_{ij} \psi_i^* \psi_j. \end{aligned}$$

First term

$$S^{(2)}[\psi^*, \psi] = \int_0^{\hbar\beta} d\tau \left\{ \sum_{ij} t_{ij} \psi_i^*(\tau) \psi_j(\tau) - \frac{1}{\hbar} \int_0^{\hbar\beta} d\tau' \sum_{iji'j'} t_{ij} t_{i'j'} \psi_j^*(\tau) \times \langle \alpha_i(\tau) \alpha_{i'}^*(\tau') \rangle_{S^{(0)}} \psi_{j'}(\tau') \right\} \quad t_{ij} = t_{ji} = \begin{cases} t & \text{for nearest neighbors} \\ 0 & \text{otherwise.} \end{cases}$$



直接带入跳跃1次的

$$\sum_{ij} t_{ij} \psi_i^*(\tau) \psi_j(\tau) = \sum_i t \psi_i^*(\tau) \psi_{i \pm \{1\}}(\tau)$$

Momentum space $\sum_{ij} t_{ij} \psi_i^*(\tau) \psi_j(\tau) = \sum_{\mathbf{k}} 2t \psi_{\mathbf{k}}(\tau) \psi_{\mathbf{k}}^*(\tau) \sum_{j=1}^d \cos(k_j a).$



• Dispersion Relations

$$\text{跳跃2次的 } \sum_{ji'j'} t_{ij}t_{i'j'} \psi_j^*(\tau) \langle a_i(\tau) a_{i'}^*(\tau') \rangle_{S^{(0)}} \psi_{j'}(\tau')$$

$$= \langle a_i(\tau) a_i^*(\tau') \rangle_{S^{(0)}} \sum_{jj'} t_{ij}t_{ij'} \psi_j^*(\tau) \psi_{j'}(\tau')$$

$$= t^2 \langle a_i(\tau) a_i^*(\tau') \rangle_{S^{(0)}} \sum_j \{ z \psi_j^*(\tau) \psi_j(\tau') + \psi_j^*(\tau) \psi_{j \pm \{2\}}(\tau') + \psi_j^*(\tau) \psi_{j \pm \{\sqrt{2}\}}(\tau') \}$$

① ② ③


$$z = 2d \quad 2 \sum_i^d \cos(2k_j a)$$

$$= \langle a_i(\tau) a_i^*(\tau') \rangle_{S^{(0)}} \sum_{\mathbf{k}} \psi_{\mathbf{k}}^*(\tau) \psi_{\mathbf{k}}(\tau') \bar{\epsilon}_{\mathbf{k}}^2, \quad \bar{\epsilon}_{\mathbf{k}} = 2t \sum_{j=1}^d \cos(k_j a).$$

$$\langle a_i^* a_j^* \rangle_{S^{(0)}} = \langle a_i a_j \rangle_{S^{(0)}} = 0,$$

$$\langle a_i^* a_j \rangle_{S^{(0)}} = \langle a_i a_j^* \rangle_{S^{(0)}} = \langle a_i a_i^* \rangle_{S^{(0)}} \delta_{i,j},$$

$$4 \sum_i^d \sum_{j \neq i}^d \cos(k_i a) \cos(k_j a)$$





• Dispersion Relations

处理时间依赖，引入Matsubara frequency $\hbar\omega_n = \pi(2n)/\hbar\beta$

$$\psi_{\mathbf{k}}(\tau) = \sum_n \frac{1}{\sqrt{\hbar\beta}} \psi_{\mathbf{k}n} e^{-i\omega_n\tau}, \quad \psi_{\mathbf{k}}^*(\tau) = \sum_n \frac{1}{\sqrt{\hbar\beta}} \psi_{\mathbf{k}n}^* e^{+i\omega_n\tau}.$$

$$E_g^{(0)} = \begin{cases} 0 & \text{if } \bar{\mu} < 0, \\ \frac{1}{2}\bar{U}g(g-1) - \bar{\mu}g & \text{if } \bar{U}(g-1) < \bar{\mu} < \bar{U}g. \end{cases}$$

引入虚的时间排序算符，将场期望转换为对算符的期望

$$\langle a_i(\tau)a_{i'}^*(\tau') \rangle_{S^{(0)}} = \langle T[a_i(\tau)a_{i'}^\dagger(\tau')] \rangle_{S^{(0)}}.$$

$$\begin{aligned} \langle a_i(\tau)a_{i'}^*(\tau') \rangle_{S^{(0)}} &= \theta(\tau - \tau') (1 + g) \\ &\times \exp\{-(E_{g+1}^{(0)} - E_g^{(0)})(\tau - \tau')/\hbar\} \\ &+ \theta(\tau' - \tau) g \exp\{(E_{g-1}^{(0)} - E_g^{(0)}) \\ &\times (\tau - \tau')/\hbar\}. \end{aligned}$$

利用阶跃函数可以表示为

$$\begin{aligned} \langle T[a_i(\tau)a_{i'}^\dagger(\tau')] \rangle_{S^{(0)}} &= \theta(\tau - \tau') \langle a_i(\tau)a_{i'}^\dagger(\tau') \rangle_{S^{(0)}} \\ &+ \theta(\tau' - \tau) \langle a_{i'}^\dagger(\tau')a_i(\tau) \rangle_{S^{(0)}}. \end{aligned}$$



$$E_{g+1}^{(0)} - E_g^{(0)} = -\mu + gU > 0,$$

$$E_g^{(0)} - E_{g-1}^{(0)} = -\mu + (g-1)U < 0.$$

e指数是考虑Heisenberg绘景导致的吗?
为什么没有i



• Dispersion Relations

代入并利用阶跃函数的性质可得

$$S^{(2)}[\psi^*, \psi] = \sum_n \sum_{\mathbf{k}} |\psi_{kn}|^2 \bar{\epsilon}_{\mathbf{k}} \times \left(1 - \frac{\bar{\epsilon}_{\mathbf{k}}}{\hbar} \int_{-\infty}^0 d\tau' (1+g) \times \exp\{(-i\hbar\omega_n - \mu + gU)\tau'/\hbar\} - \frac{\bar{\epsilon}_{\mathbf{k}}}{\hbar} \int_0^\infty d\tau' g \exp\{-(i\hbar\omega_n + \mu - (g-1)U)\tau'/\hbar\} \right)$$

即

$$S^{(2)}[\psi^*, \psi] = \sum_n \sum_{\mathbf{k}} |\psi_{kn}|^2 \bar{\epsilon}_{\mathbf{k}} \times \left[1 - \bar{\epsilon}_{\mathbf{k}} \left(\frac{g+1}{-i\hbar\omega_n - \mu + gU} + \frac{g}{i\hbar\omega_n + \mu - (g-1)U} \right) \right].$$

带入实能量 $\hbar\omega$, $i\omega_n \rightarrow \omega$, 并令其等于0, 得到等式


$$0 = \left[1 - \bar{\epsilon}_{\mathbf{k}} \left(\frac{g+1}{-\hbar\omega - \mu + gU} + \frac{g}{\hbar\omega + \mu - (g-1)U} \right) \right]$$

即为准粒子和准空穴色散关系

$$\hbar\omega_{qp,qh} = -\mu + \frac{U}{2}(2g-1) - \frac{\bar{\epsilon}_{\mathbf{k}}}{2} \pm \frac{1}{2} \sqrt{\bar{\epsilon}_{\mathbf{k}}^2 - (4g+2)U\bar{\epsilon}_{\mathbf{k}} + U^2}.$$

$$\lim_{U \rightarrow \infty} \hbar\omega_{qp} = -\mu + gU - (g+1)\bar{\epsilon}_0 \quad \lim_{U \rightarrow \infty} \hbar\omega_{qh} = -\mu + (g-1)U + g\bar{\epsilon}_0$$

$$2024/11/25 \quad = E_{g+1}^{(0)} - E_g^{(0)} - (g+1)zt, \quad = E_g^{(0)} - E_{g-1}^{(0)} + gzt,$$



一阶修正来自于hopping, i节点准粒子激发则有 $\langle c_j^\dagger c_i \rangle = (g+1)t$, 而空穴激发则 $\langle c_i^\dagger c_j \rangle = gt$